Cancer Research Consortium of West Michigan P: 616.391.1230 | 800.336.6299 | www.crcwm.org

# FAST FACTS

### NRG CC005 - FORTE (Five or Ten Year Colonoscopy for 1-2 Non-Advanced Adenomatous Polyps)

#### Eligibility Criteria

- 1. The participant must have signed and dated an IRB-approved consent form that conforms to federal and institutional guidelines.
- **2.** Participants  $\geq$  45 and < 70 years of age at the time of qualifying colonoscopy.
- **3.** Participants with a first-time diagnosis of 1-2 non-advanced tubular adenomas (<sup>□</sup> 10 mm without tubulovillous or villous changes or high grade or severe dysplasia) *from the qualifying colonoscopy* within 4 years prior to randomization.
  - a. Sessile serrated polyps/adenomas, as long as they do not meet the criteria for advanced adenomas, will be considered as non-advanced adenomas.
- **4.** Qualifying colonoscopy must be a complete colonoscopy with visualization of the cecum and with adequate cleansing within 4 years prior to randomization.
- 5. Complete excision of all observed polyps in qualifying colonoscopy (see Section 3.3.7).
- 6. Participants must be able to read or understand English or Spanish.

#### **Ineligibility Criteria**

- 1. Prior history of colorectal cancer or colorectal adenomas including sessile serrated polyps/adenomas excluding those found on the qualifying colonoscopy.
- 2. Prior history of a hyperplastic polyp measuring  $\geq$  1 cm in size.
- 3. Traditional serrated adenomas found on the qualifying colonoscopy.
- 4. Hyperplastic polyp measuring  $\geq$  1 cm in size found on the qualifying colonoscopy.
- 5. Previous malignancies unless the participant has been disease-free for 5 or more years prior to randomization and is deemed by the physician to be at low risk for recurrence. Participants with the following cancers are eligible if diagnosed and treated within the past 5 years: all in situ cancers and basal cell and squamous cell carcinoma of the skin.
- 6. Colonoscopy performed *after* the qualifying colonoscopy but prior to randomization.
- 7. Incomplete qualifying colonoscopy (e.g., cecum not visualized).
- 8. Incomplete endoscopic excision of adenomatous polyps based on colonoscopist impression at qualifying colonoscopy. (Excision of all hyperplastic rectosigmoid polyps is not required.)
- 9. Sub-total colectomy or total proctocolectomy. (Segmental resections are allowed.)
- 10. Known family history of CRC diagnosed at 2 60 years of age in a first degree relative (mother, father, child, sibling) or in two first degree relatives with CRC at any age.
- 11. Participants with a clinical diagnosis of a significant heritable risk for colorectal cancer (Familial Adenomatous Polyposis, Hereditary Nonpolyposis Colorectal Cancer [Lynch Syndrome]).
- 12. Participants tested positive for a Familial Adenomatous Polyposis, Hereditary Nonpolyposis Colorectal Cancer [Lynch Syndrome] genetic mutation that increases risk of colorectal cancer.
- 13. Inflammatory bowel disease (e.g., Crohn's Disease, ulcerative colitis).
- 14. Life expectancy less than 10 years due to comorbid conditions in the opinion of the investigator.
- 15. Other comorbid conditions that would prevent the participant from having colonoscopies or would prevent required follow-up

#### Figure 1. NRG-CC005 SCHEMA

Participants ≥ 45 and < 70 years with First Diagnosis of 1-2 Non-Advanced Adenomas within Prior 4 years

## STRATIFICATION

- Age at qualifying colonoscopy (45-55, 56 < 70)
- Sex (Female, Male)
- Time from Qualifying Colonoscopy to Randomization (<2 years, 2-4 years)



\* Randomization is 1:1.